

**Table 1.** Quality of life in patients with LLR.

NUMBER OF PATIENTS	186
AGE (years) mean $\pm$ SD	43.87 $\pm$ 14.99
SEX -Female: n (%) -Male: n (%)	96 (51.6%) 90 (48.4%)
PLACE OF RESIDENCE -Urban: n (%) -Rural: n (%)	102 (54.8%) 84 (45.2%)
INSECT IMPLICATED IN THE REACTION -Wasp: n (%) -Bee: n (%) -Unidentified: n (%)	153 (82.3%) 20 (10.8%) 13 (7%)
TYPE OF REACTION -Immediate (< 1 hour after Sting) -Non immediate (> 1 hour after Sting)	105 (56.5%) 79 (42.5%)
LOCATION OF THE REACTION -Extremities -Head and neck	151 (81.2%) 32 (17.2%)
DURATION OF THE REACTION (median) (ICR) (days)	3 (2-5)
AVERAGE SCORE HiCaVi $\pm$ SD*	4.1 (1.53)
SCORE HiCaVi BY SEX -* female -* male	3.93 (1.56); p=0.124 4.28 (1.50)
SCORE HiCaVi BY INSECT -* wasp -* bee	4.04 (1.52); p=0.054 4.74 (1.51)
SCORE HiCaVi BY TYPE OF REACTION -* immediate -*non immediate	3.87 (1.49); p=0.031 4.36 (1.56)
SCORE HiCaVi BY PLACE OF RESIDENCE -* urban -* rural	4.12 (1.37); p=0.873 4.08 (1.66)
SCORE HiCaVi BY LOCATION OF THE REACTION -* extremities -* head and neck	4.06 (1.54); p=0.756 4.15 (1.54)
SCORE HiCaVi LLR AND SR -* LLR -* SR	4.10 (1.53); p=0.145 3.86 (1.55)

Abbreviations: LLR: local large reaction; SR: systemic reaction