

SUPPLEMENTARY MATERIAL

Table 1. Quality of life in patients with LLR.

NUMBER OF PATIENTS	186
AGE (years) mean \pm SD	43.87 \pm 14.99
SEX	
-Female: n (%)	96 (51.6%)
-Male: n (%)	90 (48.4%)
PLACE OF RESIDENCE	
-Urban: n (%)	102 (54.8%)
-Rural: n (%)	84 (45.2%)
INSECT IMPLICATED IN THE REACTION	
-Wasp: n (%)	153 (82.3%)
-Bee: n (%)	20 (10.8%)
-Unidentified: n (%)	13 (7%)
TYPE OF REACTION	
-Immediate (< 1 hour after Sting)	105 (56.5%)
-Non immediate (> 1 hour after Sting)	79 (42.5%)
LOCATION OF THE REACTION	
-Extremities	151 (81.2%)
-Head and neck	32 (17.2%)
DURATION OF THE REACTION (median) (ICR) (days)	3 (2-5)
AVERAGE SCORE HiCaVi \pm SD*	4.1 (1.53)
SCORE HiCaVi BY SEX	
-* female	3.93 (1.56); p=0.124
-* male	4.28 (1.50)
SCORE HiCaVi BY INSECT	
-* wasp	4.04 (1.52); p=0.054
-* bee	4.74 (1.51)
SCORE HiCaVi BY TYPE OF REACTION	
-* immediate	3.87 (1.49); p=0.031
-* non immediate	4.36 (1.56)
SCORE HiCaVi BY PLACE OF RESIDENCE	
-* urban	4.12 (1.37); p=0.873
-* rural	4.08 (1.66)
SCORE HiCaVi BY LOCATION OF THE REACTION	
-* extremities	4.06 (1.54); p=0.756
-* head and neck	4.15 (1.54)
SCORE HiCaVi LLR AND SR	
-* LLR	4.10 (1.53); p=0.145
-* SR	3.86 (1.55)

Abbreviations: LLR: local large reaction; SR: systemic reaction