

MATERIAL SUPPLEMENTARY

1. Kourani E, Corazza F, Michel O, Doyen V. What Do We Know About Fish Allergy at the End of the Decade? *J Investig Allergol Clin Immunol*. 2019;29(6):414-21.
2. Grabenhenrich LB, Dölle S, Moneret-Vautrin A, Köhli A, Lange L, Spindler T, et al. Anaphylaxis in children and adolescents: The European Anaphylaxis Registry. *J Allergy Clin Immunol*. 2016;137(4):1128-37.e1.
3. Van Do T, Hordvik I, Endresen C, Elsayed S. Characterization of parvalbumin, the major allergen in Alaska pollack, and comparison with codfish Allergen M. *Molecular immunology*. 2005;42(3):345-53.
4. Kuehn A, Hilger C, Lehnert-Weber C, Codreanu-Morel F, Morisset M, Metz-Favre C, et al. Identification of enolases and aldolases as important fish allergens in cod, salmon and tuna: component resolved diagnosis using parvalbumin and the new allergens. *Clin Exp Allergy*. 2013;43(7):811-22.
5. Carvalho S, Marcelino J, Cabral Duarte MF, Costa C, Barbosa MA, Pereira Dos Santos MC. Role of Recombinant Parvalbumin Gad c 1 in the Diagnosis and Prognosis of Fish Allergy. *J Investig Allergol Clin Immunol*. 2020;30(5):340-5.
6. Asero R, Mistrello G, Roncarolo D, Casarini M, Falagiani P. True monosensitivity to a tropical sole. *Allergy*. 1999;54(11):1228-9.
7. Ebo DG, Kuehn A, Bridts CH, Hilger C, Hentges F, Stevens WJ. Monosensitivity to pangasius and tilapia caused by allergens other than parvalbumin. *J Investig Allergol Clin Immunol*. 2010;20(1):84-8.
8. Kuehn A, Hutt-Kempf E, Hilger C, Hentges F. Clinical monosensitivity to salmonid fish linked to specific IgE-epitopes on salmon and trout beta-parvalbumins. *Allergy*. 2011;66(2):299-301.
9. Kuehn A, Codreanu-Morel F, Lehnert-Weber C, Doyen V, Gomez-Andre SA, Bienvenu F, et al. Cross-reactivity to fish and chicken meat - a new clinical syndrome. *Allergy*. 2016;71(12):1772-81.
10. Hilger C, Thill L, Grigioni F, Lehnert C, Falagiani P, Ferrara A, et al. IgE antibodies of fish allergic patients cross-react with frog parvalbumin. *Allergy*. 2004;59(6):653-60.
11. Gonzalez-de-Olano D, Bartolome B, Maroto AS, Vivanco F, Pastor-Vargas C. Asthma after chicken consumption due to cross-reactivity between fish and chicken parvalbumin. *J Investig Allergol Clin Immunol*. 2012;22(3):227-8.
12. Haroun-Diaz E, Blanca-Lopez N, Vazquez de la Torre M, Ruano FJ, Somoza Alvarez ML, Labrador Horrillo M, et al. Severe anaphylaxis due to crocodile-meat allergy exhibiting wide cross-reactivity with fish allergens. *J Allergy Clin Immunol Pract*. 2017.
13. Ballardini N, Nopp A, Hamsten C, Vetander M, Melen E, Nilsson C, et al. Anaphylactic Reactions to Novel Foods: Case Report of a Child With Severe Crocodile Meat Allergy. *Pediatrics*. 2017.

14. Muraro A, Werfel T, Hoffmann-Sommergruber K, Roberts G, Beyer K, Bindslev-Jensen C, et al. EAACI food allergy and anaphylaxis guidelines: diagnosis and management of food allergy. *Allergy*. 2014;69(8):1008-25.
15. Pérez-Calderón R, Gonzalo-Garijo M, Rodríguez-Velasco FJ, Sánchez-Vega S, Bartolomé-Zavala B. Occupational respiratory allergy in peach crop workers. *Allergy*. 2017;72(10):1556-64.

Accepted Article

Table E1. Total IgE, skin prick test, and specific IgE (ImmunoCAP) to fish species

Patient	Skin prick test/ sIgE (kU _A /L)							
	Total IgE (UI/ml)	Cod	Tuna	Salmon	Hake	Sardine	Megrim	Swordfish
1	622	+/4.3	-/ND	+/1.2	+/1.52	+/0.15	+/2.01	+/0.26
2	103	-/2.16	-/ <0.1	-/0.42	+/0.52	-/0.21	-/ 0.55	-/0.11
3	542	+/179	+/9.61	+/22.5	+/60	+/11.9	+/52.1	+/1.45
4	44	+/5.43	-/ <0.10	+/0.22	+/1.34	-/ <0.1	+/0.53	-/ <0.1
5	108	+/39.3	+/1.63	+/2.04	+/ 15.8	+/0.64	+/11	+/10.6
6	238	+/2.47	-/0.24	+/1.03	+/ 0.73	+/0.24	+/0.55	-/ <0.1
7	1734	+/9.13	-/0.39	+/2.96	+/ 2.75	+/0.41	+/1.86	-/0.19
8	116	+/45.6	+/1.38	+/1.17	+/ 9.14	+/2.01	+/2.2	-/0.11
9	2367	+/100	+/2.7	+/68	+/ 15.4	+/5.4	+/8.1	-/1.3
10	252	+/30.5	+/1.76	+/15.5	+/ 5.45	+/3.43	+/3.26	-/0.5
11	129	+/21.7	+/0.65	+/6.27	+/ 3.86	+/1.11	+/2.59	-/0.19
12	495	+/10.6	+/0.98	+/2.39	+/ 3.38	+/0.74	+/4	+/1.91
13	398	ND/53.7	ND/6.47	ND/20.2	ND/13.8	ND/6.08	ND/10.5	ND/1.75
14	4396	+/94	+/4.36	+/34.6	+/ 24	+/5.56	+/17.5	-/1.55
15	6323	+/2.7	-/0.15	+/0.22	+/ 1.67	+/0.18	+/1.54	+/1.73
16	1070	+/15	+/2.89	+/12.7	+/ 4.96	+/3.1	+/7.06	-/1.61
17	2990	+/78	+/2.75	+/28.7	+/ 34.5	+/7.23	+/22.2	-/1.73
18	797	+/42.3	+2.42	+/14.9	+/ 8.06	+/2.81	+/10.8	-/0.44
19	110	+/17.4	+/3.15	+/26	+/ 3.95	+/4.64	+/3.21	+/0.92
20	9656	+/64.7	+/0.68	+/7.42	+/ 14.4	+/2.56	+/1.78	+/1.04
21	1778	+/40.2	+/1.06	+/6.67	+/ 7.43	+/1.76	+/9.68	-/1.15
22	2993	+/28.8	+/1.12	+/1.94	+/ 8.99	+/0.92	+/11.3	+/3.83
23	24	+/21.2	+/0.97	+/10.3	+/ 0.94	+/1.38	+/0.65	-/0.7
24	543	-/ <0.1	-/ <0.1	-/ <0.1	-/ <0.1	-/ <0.1	+/1.77	-/0.13
25	825	-/0.79	+/0.39	-/0.35	+/ 0.81	-/ <0.1	+/1.99	+/0.69
26	1774	+/5.05	-/0.44	+/1.54	+/ 1.18	+/0.7	+/1.63	-/0.38
27	23	+/5.11	-/0.22	+/1.48	+/ 0.74	+/0.39	+/0.78	-/ <0.1

+ : Positive. A wheal diameter larger than 3 mm was considered a positive result; - :

Negative; ND: not done.

* Positive values were those ≥ 0.1 kU_A/L

Table E2. Skin test and specific IgE to crocodile, frog, and chicken extracts

Patient	CROCODILE					FROG			CHICKEN		
	Prick test		Prick-prick		EAST kU _A /L	Prick-prick		EAST kU _A /L	Prick-prick		CAP kU _A /L
	RCrme	BCrme	RCrPP	BCrPP	Cr	RFPP	BFPP	F	RChPP	BChPP	Ch
1	ND	ND	+	+	<0.35	ND	ND	0.5	ND	ND	ND
2	ND	ND	-	-	<0.35	ND	ND	0.4	ND	ND	ND
3	+	+	+	+	15.6	+	+	15.8	+	+	<0.10
4	-	-	-	-	0.4	-	-	0.5	+	-	<0.10
5	+	+	+	+	2.3	+	+	1.3	+	+	1.02
6	-	-	-	-	<0.35	-	-	<0.35	+	-	0.24
7	+	-	+	+	0.7	-	-	<0.35	-	-	0.13
8	+	+	+	+	0.4	+	+	<0.35	+	-	<0.10
9	ND	+	+	+	7.0	ND	ND	1.7	ND	ND	<0.10
10	+	+	+	+	2.4	-	+	2.9	+	+	0.83
11	+	+	+	+	0.4	+	+	0.6	+	+	<0.10
12	+	+	+	+	2.6	+	+	1.6	+	+	2.79
13	ND	ND	ND	ND	3.0	ND	ND	3.8	ND	ND	<0.10
14	+	+	+	+	8.2	+	+	4.8	+	+	0.85
15	+	+	+	+	3.4	+	+	2.9	+	+	4.05
16	+	+	+	+	1.4	-	-	0.9	+	-	0.98
17	+	+	+	+	12	+	+	9.4	+	-	0.42
18	+	+	+	+	0.9	+	ND	1.0	-	ND	0.16
19	+	+	+	-	0.4	+	+	0.4	-	-	<0.10
20	-	-	+	-	9.0	-	-	3.8	+	-	1.76
21	+	+	+	+	2.1	+	+	5.7	+	+	1.48
22	+	+	+	+	35.6	+	+	16.2	+	+	74.7
23	-	-	-	-	0.8	-	-	<0.35	-	-	<0.10
24	+	-	+	-	0.9	+	-	0.4	+	-	0.4
25	-	-	-	-	0.4	+	-	<0.35	+	-	0.22
26	+	-	+	-	0.4	+	+	1.9	+	-	0.4
27	+	+	+	-	<0.35	+	-	<0.35	-	-	<0.10

Cr: crocodile; F: frog; Ch: chicken; RCrme: raw crocodile meat extract; BCrme: boiled crocodile meat extract; RCrPP: raw crocodile prick-prick; BCrPP: boiled crocodile prick-prick; RFPP: raw frog prick-prick; BFPP: boiled frog prick-prick; RChPP: raw chicken prick-prick; BChPP: boiled chicken prick-prick; ND: not done.

* Positive values were those ≥ 0.1 kU_A/L (Siemens Immulite 2000/Xpi®), and ≥ 0.35 kU_A/L (EAST)