

Basophil activation test with progressively less heated forms of egg distinguishes egg allergic from tolerant children

Running title: BAT with progressively less heated egg extracts

De Vlieger L¹, Nuyttens L^{1,2}, Ieven T^{1,3}, Diels M², Coorevits L¹, Cremer J¹, Schrijvers R^{1,3}, Bullens DMA^{1,2}

¹Department of Microbiology, Immunology and Transplantation, Allergy and Immunology Research Group, KU Leuven, Leuven, Belgium

²Department of Pediatrics, University Hospitals Leuven, Leuven, Belgium

³Department of General Internal Medicine, University Hospitals Leuven, Leuven, Belgium

Correspondence:

Dominique Bullens

UZ Herestraat 49 – bus 811

3000 Leuven

E-mail: dominique.bullens@kuleuven.be

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.18176/jiaci.0964

Abstract

Background: Diagnosis of egg allergy through basophil activation testing (BAT) has been mainly performed with an egg white extract or individual egg allergens rather than clinically more representative whole-egg extracts. Impact of heating on whole-egg extract allergenicity remains unassessed.

Objective: Validating BAT with gradually less heated whole-egg extracts in egg allergy diagnosis and as tolerance marker.

Methods: CD63-based BAT was performed with five progressively less heated extracts from cake, hard-boiled egg, omelet, soft-boiled, and raw egg in 10 egg allergic (EA), 10 complete egg tolerant (ET) and 12 non-egg-sensitized non-allergic (NEA) children. Cutoffs and diagnostic accuracy measures were established through ROC analysis. Changes in basophil response were assessed in 12 baked egg tolerant children undergoing an 8-month gradual egg reintroduction protocol with BAT and oral food challenges prior to each reintroduction step.

Results: Basophil responses to all egg extracts were increased in EA, but not in ET and NEA children. Responses decreased progressively with more heated egg extracts. Compared to ET children, EA children showed higher basophil sensitivity for all egg extracts. Negative BAT responses predicted clinical tolerance with a 90-100% sensitivity, 100% specificity, and false positive rate of 2.78%. In comparison, egg sIgE's (<0.35 kUA/L) had a lower specificity of 50-78% with a false positive rate of 40%. Basophil reactivity and sensitivity tended to decrease in baked egg tolerant children undergoing gradual egg reintroduction, concurrent with tolerance development.

Conclusion: BAT with progressively less heated egg preparations is a sensitive and highly specific tool to discriminate EA from ET children.

Key words: Basophil. Basophil activation test. Egg allergy. Baked egg tolerance. Heated egg. Pediatric. Egg sIgE.

Resumen

Antecedentes: El diagnóstico de la alergia al huevo mediante test de activación de basófilos (TAB) se ha realizado principalmente con un extracto de clara de huevo o alérgenos de huevo individuales en lugar de con extractos de huevo entero clínicamente más representativos. Aún no se ha evaluado el impacto del calentamiento en la alergenidad del extracto de huevo entero.

Objetivo: Validar el TAB con extractos de huevo entero gradualmente menos calentados en el diagnóstico de la alergia al huevo y como marcador de tolerancia.

Métodos: Se realizó un TAB basado en la expresión de CD63 con cinco extractos de huevo progresivamente menos calentados (pastel, huevo duro, tortilla, huevo pasado por agua y huevo crudo) en 10 niños alérgicos al huevo (AH), 10 completamente tolerantes al huevo (TH) y 12 no alérgicos ni sensibilizados al huevo (NA). Se establecieron puntos de corte y medidas de precisión diagnóstica mediante análisis ROC. Se evaluaron los cambios en el TAB en 12 niños que toleraban el huevo horneado sometidos a un protocolo de reintroducción gradual del huevo durante 8 meses con TAB y provocaciones orales previas a cada paso de la reintroducción.

Resultados: La respuesta de basófilos a todos los extractos de huevo estaban aumentadas en los niños AH, pero no en los TH y NA. Las respuestas disminuyeron progresivamente con extractos de huevo más calentados. En comparación con los niños TH, los niños AH mostraron una mayor sensibilidad de los basófilos a todos los extractos de huevo. El TAB negativo predijo tolerancia clínica con una sensibilidad del 90-100%, una especificidad del 100% y una tasa de falsos positivos del 2.78%. En comparación, la IgE específica a huevo <0.35 kUA/L tuvo una especificidad inferior del 50-78% con una tasa de falsos positivos del 40%. La reactividad y la sensibilidad de los basófilos tendieron a disminuir en los niños sometidos a la reintroducción gradual de huevo, en paralelo al desarrollo de tolerancia.

Conclusión: El TAB con preparados de huevo progresivamente menos calentados es una herramienta sensible y altamente específica para discriminar a los niños alérgicos a huevo de los tolerantes.

Palabras clave: Basófilo. Test de activación de basófilos. Alergia a huevo. Tolerancia a huevo horneado. Huevo calentado. Pediátrico. IgE específica a huevo.

Summary Box

- **What do we know about this topic?**

Diagnosis of egg allergy through basophil activation testing (BAT) has been mainly performed with an egg white extract or individual egg allergens rather than clinically more representative whole-egg extracts. The impact of heating on whole-egg extract allergenicity remains unassessed.

- **How does this study impact our current understanding and/or clinical management of this topic?**

BAT with gradually less heated whole-egg extracts discriminates egg allergic from tolerant children, with superior specificity compared to egg sIgE < 0.35 kUA/L. Progressive heating reduces the ability of whole-egg extracts to induce basophil activation *in vitro*.

Introduction

Diagnosis of childhood egg allergy is currently established by medical history accompanied by a first-line diagnostic test including measurement of serum specific IgE (sIgE) or skin prick testing (SPT) [1,2]. The specificity of these first-line tests can be suboptimal as they often reflect irrelevant sensitization to hen's egg rather than clinical hen's egg allergy [1,2]. In a 2014 meta-analysis, egg sIgE levels ≥ 0.35 kUA/L predicted egg reactivity with a mean specificity of 49% and sensitivity of 93% [3]. Increasing cutoff levels offered increased specificity at the cost of sensitivity [4]. Consequently, an in-hospital oral food challenge (OFC) is the gold standard to confirm the diagnosis of egg allergy or monitor for allergy resolution [5,6]. However, this procedure requires an experienced clinical team as well as well-equipped facilities as life-threatening allergic reactions during OFCs have been described [5,6]. In this regard, basophil

activation testing (BAT) has emerged as an alternative non-invasive *ex vivo* assay for IgE-mediated hypersensitivity that can be used to diagnose food allergy and monitor the development of natural or immunotherapy-induced tolerance [7–9].

The majority of studies that assessed the performance of BAT to diagnose egg allergy or monitor for resolution used an egg white extract or the individual egg allergens, ovalbumin and ovomucoid, for basophil stimulation [10–12]. An ovalbumin-based BAT could diagnose egg allergy with a 100% specificity and 77% sensitivity [10]. In another study, basophil reactivity following stimulation with egg white extract discriminated two clinical egg allergy phenotypes, in which baked egg reactive children had a higher percentage of CD63-positive (%CD63⁺) basophils to egg white compared to baked egg tolerant (BET) children [13]. Several studies evaluating BAT during egg oral immunotherapy also reported a decreased %CD63⁺ basophils to pasteurized whole egg, egg white, ovalbumin or ovomucoid at the end of the treatment [14–19]. However, few if any studies examined BAT with clinically representative whole-egg extracts and no studies have evaluated BAT as a non-invasive predictor of clinical tolerance during the gradual reintroduction of egg using an egg ladder [14,18].

In a recent study, we demonstrated the safe induction of tolerance to raw egg in a BET cohort through progressive introduction of less heated egg products over a 24-month period [20]. Having characterized extracts from these egg products (cake, hard-boiled egg, omelet, soft-boiled egg and raw egg), we first aimed to investigate the diagnostic performance of BAT with these extracts in discriminating between true egg allergic and tolerant children with positive SPT or sIgE to egg or egg components. Secondly, we assessed the evolution and predictive value of basophil responses in BET children undergoing a shortened 8-month gradual egg reintroduction protocol.

Methods

Basophil activation testing

BAT was performed on fresh heparinized whole blood samples (Suppl. Material for full protocol). In brief, samples were stimulated for 20 minutes at 37°C with stimuli dissolved in an IL3-containing buffer (final concentration 9 ng/mL). Stimuli included 10-fold serial dilutions of five progressively less heated whole-egg extracts (0.1-100 µg/mL) including cake (35' at 165°C), hard-boiled egg (10' at 100°C), omelet (4' at 120°C), soft-boiled egg (5' at 100°C) and raw egg. Extracts were prepared as previously described, and characterized in our previously published work (Suppl. Material) [20]. Mono- and polyclonal anti-human IgE (aIgE, 5 µg/mL), formyl-methionyl-leucyl-phenylalanine (fMLP, 2 µM) or buffer alone were used as positive and negative controls, respectively. Stimulation was halted on ice followed by staining with anti-CD123 PE, anti-HLA-DR AF647 and anti-CD63 FITC. After erythrocyte lysis, a minimum of 500 basophils ($SSC^{low}/CD123^{+}/HLA-DR^{-}$) were acquired on the LSR Fortessa flow cytometer running DIVA software, and analysed with FlowJo 10.8.1. Basophil activation was measured as %CD63⁺ basophils, corrected for spontaneous CD63-expression by subtracting the %CD63⁺ basophils in the unstimulated control condition. Children with < 5 %CD63⁺ basophils to mono- and polyclonal aIgE were classified as non-releasers, henceforth termed non-responders [9].

Diagnostic cohort

To explore the discriminative capacity of BAT between true egg allergy and tolerance, we recruited 10 egg allergic (EA), 10 egg-sensitized but tolerant (ET) and 12 non-egg-sensitized non-allergic (NEA) children from the Pediatrics department of UZ Leuven. Egg allergy was

defined as a clinical type I hypersensitivity response to egg along with a positive egg SPT (wheal > 3 mm) or egg sIgE \geq 0.35 kUA/L (ImmunoCAP, lower limit of quantification of 0.10 kUA/L). Children with a prior history of egg allergy who ingested foods containing raw egg without developing symptoms at the time of inclusion, independent of their egg sIgE's, were considered complete egg tolerant.

Gradual egg reintroduction cohort (pre-TETI-II study)

Changes in BAT outcome were evaluated in an ongoing pilot study (pre-TETI-II) involving 12 BET children (L1-L12) consecutively reintroducing cake, hard-boiled egg, omelet, soft-boiled and raw egg over a period of 8 months [20] (Figure 1). This 8-month time period was supported by findings from our previous study, as several children progressed through the step-wise protocol at accelerated pace (parental decision) and safely developed raw egg tolerance within 5-12 months [20]. Included children had proven egg allergy along with an ovomucoid sIgE predicting at least a 75% chance for passing baked egg OFC, but were still supposed to react to less heated egg products [21,22]. Prior to each reintroduction step, children underwent an in-hospital OFC with the corresponding egg preparation during which an additional blood sample was collected for BAT with all five egg extracts (Table S1). If OFCs were tolerated, cake and hard-boiled egg were further introduced for 3 months at home followed by a 1 month introduction of omelet and soft-boiled egg. Both studies were approved by the Ethics Committee Research UZ/KU Leuven and written informed consent was obtained from parents with accompanying assent of the child from the age of six years onwards.

Statistics

Statistical analysis was performed using GraphPad Prism v.9.2.0 for Windows. Normality was determined using the D'Agostino and Pearson test. Continuous variables are reported as

medians (interquartile range) or means (95% confidence interval) and compared between groups using the student t-test or mixed-model analysis when appropriate. Cubic spline and non-linear regression analysis were used to model the basophil CD63 dose-response for each extract. Receiver operator characteristic (ROC) curves were constructed to compare the area under the curve (AUC) for each extract concentration and to determine optimal cutoffs for BAT positivity based on optimal sensitivity and specificity values. Concentrations eliciting half-maximal basophil activation (EC50) were derived from best-fit dose-response curves for each extract. Allergen threshold sensitivity (CD-sens) was calculated by the formula “ $1/EC50 \times 100$ ” [9]. Correlations between egg sIgE’s and BAT responses were evaluated through the Spearman or Pearson rank correlation test where appropriate. BAT non-responders were excluded from the statistical analysis of all BAT data. A *p*-value below 0.05 was considered statistically significant.

Results

Discriminative capacity of the BAT between egg allergy and tolerance

Among the 10 EA, 10 ET, and 12 NEA children, 2 (6.3%) had non-responder basophils to aIgE and were excluded from further analysis. Characteristics of included ET and EA children are depicted in Table 1. Basophils of EA children showed high CD63-expression when stimulated with the five egg extracts, with the mean %CD63⁺ basophils increasing in response to progressively less heated forms of egg (i.e. increasingly allergenic) (Figure 2, Table S2). In contrast, basophils of ET and NEA children showed low CD63-expression upon stimulation with these five egg extracts, with no significant difference between both groups.

Basophil dose-responses of EA children differed across the five extracts, with cake inducing a progressive increase in CD63-expression up to the maximum concentration of 100 µg/mL whereas reactivity to hard-boiled egg reached a plateau at 0.10 µg/mL. Omelet, soft-boiled egg and raw egg induced bell-shaped dose-responses with a decreasing CD63-expression from 1 µg/mL onwards (Figure 2). Compared to cake, the %CD63⁺ basophils was significantly higher in EA children when stimulated with 0.1 µg/mL of hard-boiled egg, omelet and soft-boiled egg, and 1 µg/mL of omelet and raw egg (Figure S1). CD63-expression differed significantly between EA and ET children across all concentrations for all tested extracts with the exception of 0.1 µg/mL of cake, which could not distinguish EA from ET children (Figure S2). The AUC of the BAT with each egg extract also significantly differed between EA and ET children (Figure S3). Accordingly, EA children showed higher basophil sensitivity for all egg extracts, as expressed by a lower EC50 compared to ET children (Figure 2).

Optimal %CD63⁺ cutoff values for discrimination between EA and ET children were calculated for all tested concentrations (0.1-100 µg/mL), and all five egg extracts using ROC analysis (Table 2). The area under the ROC curve (AUC ROC) ranged between 0.98 and 1, indicating excellent discriminative capacity between both groups. Using these cutoffs, BAT sensitivity in discriminating EA from ET children ranged from 90% to 100% with a specificity of 100% and false positive rate (FPR) of 2.78% (data not shown). Only the 0.1 µg/mL cake extract had a lower AUC ROC of 0.65, corresponding to a 66.67% sensitivity and 55.56% specificity. When comparing this to the diagnostic accuracy of egg sIgE's using the classic cutoff of 0.35 kUA/L, the AUC ROC ranged from 0.89 to 1, with corresponding sensitivities between 90-100%, and specificities between 50-78% (Table S3). Ovomuroid sIgE had the highest AUC ROC value (1.00), which could discriminate EA from ET children with a 100% sensitivity and 60% specificity, resulting in a FPR of 40%. Alternatively, using the lower limit of detection of 0.10 kU/L for egg sIgE's resulted in an even lower specificity between 20-44% (Table S3). Next, we sought to improve diagnostic accuracy of egg sIgE's by calculating optimal cutoffs for each sIgE based on ROC analysis (Table S3). When applying these cutoffs in our cohort, higher specificities ranging from 80 to 100 % were achieved with minimal reduction in sensitivity. Additionally, egg white and ovalbumin sIgE's correlated positively with BAT dose-response AUC and maximal %CD63⁺ basophils in response to cake, hard-boiled egg, and omelet (Table S4). Ovomuroid sIgE's also positively correlated with the AUC of cake ($r=0.51$, $p=0.03$).

Outcome of gradual egg reintroduction and comparison of OFC and BAT

Out of the 12 included BET children, 9 successfully passed all five OFCs and subsequent home-based introduction without major symptoms (Table S5). Two children stopped the study earlier due to an allergic reaction during the hard-boiled egg (L8: grade I) or soft-boiled egg (L2: grade II) OFC. No clinically significant changes in blood pressure, tryptase or complement components were observed during OFCs in any of the children (Table S6). Patient L9 and L11 had elevated tryptase levels before OFCs, which was most likely attributable to hereditary alpha-tryptasemia given absence of signs of primary mast cell disease. Though, genetic testing was not available at that time. Based on current evidence, no interaction with food allergy nor basophil responses was expected [24]. One child (L3) discontinued the study due to an itchy tongue whenever eating runny egg yolk at home, despite passing the soft-boiled egg OFC. The clinical features of the study population are provided in Table 3.

To evaluate whether BAT could predict OFC-associated symptoms, we applied the previously defined %CD63⁺ cutoff values at 10 µg/mL of each extract, which had the highest sensitivity and specificity as well as the lowest workload (fewer dilutions), thus decreasing the risk of errors by manipulation (Table 2). Table 4 shows BAT responses to each extract with the outcome of their corresponding OFC. Three children (L8, 10, 11) were non-responders at the start, with one child (L11) becoming a responder at visit (V) three. Overall, out of 45 study visits with available results for concurrent BAT and OFC, nine (20%) were uninterpretable due to non-responding basophils. From the remaining 36 informative BATs, 31 (86.1%) were concordant with OFC outcome whereas 5 (13.8%) were false positive.

Evolution of BAT responses during tolerance development

No significant changes were noted in the %CD63⁺ basophils to aIgE and fMLP from baseline to end of the study, including those with non-responding basophils (Figure S4). Overall, we observed a decreasing trend in basophil responses to all five egg extracts between the first and last study visit, though these differences did not reach statistical significance (Figure 3, Figure S5). This decrease in the %CD63⁺ basophils was most pronounced in the BAT with omelet followed by soft-boiled egg, hard-boiled egg, raw egg and cake, respectively (Figure 4). Additionally, basophil sensitivity decreased for all egg extracts over the course of the graduated protocol, as expressed by an increasing EC50 from first to last study visit (Figure 3). This increase in EC50 was most pronounced for raw egg, while only a small difference was found for cake. No significant changes were noted in egg sIgE's and total IgE from baseline to end of the study (Figure S6). We found a positive correlation between egg white and ovalbumin sIgE's and the dose-response AUC and maximal %CD63⁺ basophils to cake at the first study visit. Similarly, at the final study visit, the dose-response AUC of the BAT with soft-boiled egg correlated positively with egg white sIgE's ($r=0.69$, $p=0.04$, Table S7).

Discussion

In this study, BAT was evaluated as a diagnostic tool for egg allergy using progressively less heated forms of egg. We demonstrate that our BAT protocol can discriminate clinically relevant from irrelevant IgE sensitization in egg allergic versus complete egg tolerant children, with a lower FPR and superior specificity to egg sIgE's applying the classic 0.35 kUA/L cutoff.

Until now, several studies have evaluated the diagnostic accuracy of BAT to egg white or native egg proteins such as ovalbumin and ovomucoid [10–12]. Our study is the first to validate BAT with whole-egg extracts prepared under different heating conditions, which more closely approximate egg exposure in daily life. In our previous study, we characterized these extracts and demonstrated that heating led to the disappearance of ovalbumin due to formation of insoluble aggregates [20]. Recently, Claude *et al.* showed that heat-aggregated ovalbumin had a lower basophil degranulation ability than native ovalbumin [25]. In our group of EA children, basophil reactivity was also higher to raw egg containing native ovalbumin and decreased in response to variously heated forms of egg containing heat-aggregated ovalbumin. Hereby, basophil responses to cake were significantly lower compared to several less heated egg extracts at 0.1 and 1 $\mu\text{g/mL}$. Although, no significant difference was found between basophil responses to hard-boiled egg, omelet, soft-boiled egg and raw egg, despite the difference in thermal processing and consequently allergenicity. This could indicate that these EA children weren't close to acquiring tolerance to these less heated egg forms. Additionally, basophil reactivity at the maximal concentration of cake was higher compared to the degranulation observed at similar concentrations of more allergenic forms of egg. We hypothesize that at higher concentrations of cake, digestion becomes a more important factor in this extract-based set-up. Indeed, wheat further decreases the allergenicity of egg proteins by hampering their accessibility to digestion *in vivo*, which was not accounted for in our experimental setup [26].

In comparison to the classic sIgE cutoff of 0.35 kUA/L, BAT predicted egg reactivity with a superior specificity and lower FPR. Lowering the cut-off to 0.10 kUA/L for analysing the results offered an even lower specificity compared to 0.35 kUA/L, but the clinical relevance of sIgE's between 0.10-0.35 kUA/L still remains a matter of debate for certain allergens like hen's

egg [27]. However, when optimal cutoffs were selected from the ROC curve equivalent specificities to BAT were achieved with minimal loss in sensitivity. This shows that clinical decision points of egg sIgE's largely depend on the study population with age, atopic comorbidities and severity of the allergic reaction being influencing factors [28]. Ultimately, demonstrating that a uniform cutoff of 0.35 kUA/L has limited clinical relevance, although it remains widely applied in clinical practice. The optimal cutoffs for egg sIgE's and BAT in this study should therefore also be validated in a larger cohort of egg allergic and tolerant children.

In the pre-TETI-II study, 9 out of 12 BET children developed complete raw egg tolerance within approximately 8 months while 3 children experienced adverse events leading to early withdrawal. Two allergic reactions took place during the hard-boiled and soft-boiled egg OFC while one child reacted during home-based introduction of soft-boiled egg despite passing the corresponding OFC. Overall, BAT and OFC outcomes were concordant in over 86% of informative cases. Nevertheless, the small number of children experiencing allergic reactions during OFCs, and the absence of interpretable concurrent BAT results for the few in-hospital reactors preclude us from drawing a definitive conclusion on the ability of BAT to predict OFC-associated symptoms. It must be noted that two children passed OFCs, while BAT with the corresponding egg extract was positive. We cannot rule out that these children might have reacted during the OFC if a higher dose would have been administered (e.g. 4.4 g of egg white protein as per EAACI guidelines vs 2.82 g; Figure 1) [29]. Indeed, in subject L2, the positive BAT at V2 might have been an early indication of decreased tolerance which predisposed for the positive OFC at V4.

During the course of our gradual reintroduction protocol, basophil reactivity and sensitivity tended to decrease to all five egg preparations, though these differences did not reach statistical

significance. Previous egg oral immunotherapy studies showed significant decreases in CD63-expression to egg (white), ovalbumin or ovomucoid along with clinical evidence of tolerance development at the end of the treatment [14–19]. Possible explanations for our lack of observed significant difference in basophil reactivity includes small sample size, relatively high percentage of non-responders and short duration of the gradual protocol. Indeed, studies have shown that basophil reactivity can be influenced by the duration of immunotherapy as well as the dose of the food allergen [30]. The higher proportion of non-responders in our egg reintroduction (20%, 9/45 visits) and diagnostic cohort (6%, 2/32 children), compared to 15% stated in literature, was likely due to coincidence given the limited sample size [9]. Additionally, we included BET children with a transient egg allergy phenotype, who already had lower basophil reactivity to the five whole-egg extracts at baseline compared to EA children, despite only tolerating baked egg. Though, these EA children were still far from acquiring baked egg tolerance. On the other hand, Kim *et al.* also found no significant decrease in %CD63⁺ basophils to egg white after treatment with muffin over a 2-year period [17].

Clearly, validation of BAT as a marker of tolerance induction would require a larger cohort of BET children undergoing gradual reintroduction over a longer period of time. To this end, we are currently studying the evolution of the CD63-based BAT to gradually less heated egg preparations in a larger multicentric cohort of BET children undergoing a 12- or 20-month gradual egg-introducing protocol (TETI-II study, NCT04677790). A limitation of our study was reliance on a CD63/IL-3 based protocol with omission of CD203c as an additional activation marker, since IL-3 upregulates CD203c in an allergen-independent manner, limiting its interpretability [9,30,31]. Lastly, we didn't evaluate the influence of natural egg-tolerance development on the BAT of EA children after 8 months, which could have influenced the BAT

of BET children undergoing the 8-month protocol. Additional limitations to consider when implementing BAT in clinical practice are the prevalence of non-responders, inevitably resulting in uninterpretable BAT results, as well as the need for fresh blood, trained personnel, a flow cytometer and standardization (protocol, extracts) [9,31].

Lastly, we found a significant correlation between egg white and ovalbumin sIgE's and basophil responses to cake, hard-boiled egg and omelet in EA and ET children. This is concurrent with earlier observations by Kim *et al.* who found that the %CD63⁺ basophils to egg white positively correlated with egg white sIgE's of children with and without egg allergy [12]. Additionally, BAT dose-response AUC to cake correlated with ovomucoid sIgE's, which is to be expected as low IgE's to heat stable ovomucoid have been associated with a higher chance of tolerating baked egg [21,22]. Similarly, in the gradual reintroduction cohort, egg white sIgE's and basophil response to either cake or soft-boiled egg correlated positively at the beginning and end of the treatment, respectively. This evolution could be seen as a reflection of development of clinical tolerance in EA children which starts with tolerance to baked egg and ends with tolerance to lightly cooked and raw egg. In the future, it could be of interest to evaluate integration of BAT results with egg sIgE's in a larger cohort of children undergoing OFCs to construct a predictive model for tolerance development. From a diagnostic standpoint, similar to the approach proposed by Santos *et al.* in peanut allergy, use of BAT as a second-line diagnostic tool after sIgE measurement could be an accurate and cost-efficient diagnostic method in hen's egg allergy [23].

In conclusion, we demonstrate for the first time that BAT with progressively less heated egg extracts is a sensitive and highly specific tool to discriminate egg allergic from egg-sensitized children who have completely outgrown their egg allergy. In the future, it would be interesting

to compare these whole-egg extracts with classical egg white extracts and individual egg allergens to determine which strategy offers optimal discriminative capabilities. Egg sIgE measurement remains a valuable first-line diagnostic tool. However, allergen- and patient-specific cutoffs are required to optimize diagnostic accuracy in distinguishing between sensitization and true allergy. As BET children evolved along the gradual process of tolerance development, basophil reactivity and sensitivity to progressively less heated forms of egg tended to decrease over time while tolerance was installed. Additional studies in larger cohorts of BET children undergoing gradual introduction over longer time periods are ongoing to help determine the value of BAT with whole-egg extracts as a non-invasive tool to predict clinical outcome and tolerance induction.

Congress

This work was presented at the EAACI Food Allergy School (2023, Taormina) and awarded with a price for best oral presentation.

Funding statement

DB and RS are recipients of a senior clinical investigator fellowship from the Fund for Scientific Research Flanders (FWO; 1801019N 1805518N, and 1805523N). The study was supported with a TBM project grant from FWO (T001419N) and KOOR funding from UZ Leuven.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Oriel RC, Wang J. Diagnosis and Management of Food Allergy. *Pediatr Clin North Am*. 2019 Oct;66(5):941–54.
2. Flores Kim J, McCleary N, Nwaru BI, Stoddart A, Sheikh A. Diagnostic accuracy, risk assessment, and cost-effectiveness of component-resolved diagnostics for food allergy: A systematic review. *Allergy*. 2018 Aug;73(8):1609–21.
3. Soares-Weiser K, Takwoingi Y, Panesar SS, Muraro A, Werfel T, Hoffmann-Sommergruber K, et al. The diagnosis of food allergy: a systematic review and meta-analysis. *Allergy*. 2014 Jan;69(1):76–86.
4. Sampson HA. Utility of food-specific IgE concentrations in predicting symptomatic food allergy. *J Allergy Clin Immunol*. 2001 May;107(5):891–6.
5. Bird JA, Leonard S, Groetch M, Assa'ad A, Cianferoni A, Clark A, et al. Conducting an Oral Food Challenge: An Update to the 2009 Adverse Reactions to Foods Committee Work Group Report. *J Allergy Clin Immunol Pract*. 2020 Jan;8(1):75-90.e17.
6. Perry TT, Matsui EC, Conover-Walker MK, Wood RA. Risk of oral food challenges. *J Allergy Clin Immunol*. 2004 Nov;114(5):1164–8.
7. Santos AF, Shreffler WG. Road map for the clinical application of the basophil activation test in food allergy. *Clin Exp Allergy*. 2017 Sep;47(9):1115–24.
8. Briceno Noriega D, Teodorowicz M, Savelkoul H, Ruinemans-Koerts J. The Basophil Activation Test for Clinical Management of Food Allergies: Recent Advances and Future Directions. *J Asthma Allergy*. 2021 Nov;Volume 14:1335–48.
9. Santos AF, Alpan O, Hoffmann H. Basophil activation test: Mechanisms and considerations for use in clinical trials and clinical practice. *Allergy*. 2021 Aug 27;76(8):2420–32.
10. Ocmant A, Mulier S, Hanssens L, Goldman M, Casimir G, Mascart F, et al. Basophil activation tests for the diagnosis of food allergy in children. *Clin Exp Allergy*. 2009 Aug;39(8):1234–45.
11. Sato S, Tachimoto H, Shukuya A, Kurosaka N, Yanagida N, Utsunomiya T, et al. Basophil activation marker CD203c is useful in the diagnosis of hen's egg and cow's milk allergies in children. *Int Arch Allergy Immunol*. 2010;152(SUPPL. 1):54–61.
12. Kim YH, Kim YS, Park Y, Kim SY, Kim KW, Kim HS, et al. Investigation of basophil activation test for diagnosing milk and egg allergy in younger children. *J Clin Med*. 2020;9(12):1–12.
13. Berin MC, Grishin A, Masilamani M, Leung DYM, Sicherer SH, Jones SM, et al. Egg-specific IgE and basophil activation but not egg-specific T-cell counts correlate with

- phenotypes of clinical egg allergy. *J Allergy Clin Immunol*. 2018 Jul;142(1):149-158.e8.
14. Giavi S, Vissers YM, Muraro A, Lauener R, Konstantinopoulos AP, Mercenier A, et al. Oral immunotherapy with low allergenic hydrolysed egg in egg allergic children. *Allergy*. 2016 Nov;71(11):1575–84.
 15. Vila L, Moreno A, Gamboa PM, Martínez-Aranguren R, Sanz ML. Decrease in antigen-specific CD63 basophil expression is associated with the development of tolerance to egg by SOTI in children. *Pediatr Allergy Immunol*. 2013;
 16. Itoh-Nagato N, Inoue Y, Nagao M, Fujisawa T, Shimojo N, Iwata T, et al. Desensitization to a whole egg by rush oral immunotherapy improves the quality of life of guardians: A multicenter, randomized, parallel-group, delayed-start design study. *Allergol Int*. 2018 Apr;67(2):209–16.
 17. Kim EH, Perry TT, Wood RA, Leung DYM, Berin MC, Burks AW, et al. Induction of sustained unresponsiveness after egg oral immunotherapy compared to baked egg therapy in children with egg allergy. *J Allergy Clin Immunol*. 2020 Oct;146(4):851-862.e10.
 18. Burks AW, Jones SM, Wood RA, Fleischer DM, Sicherer SH, Lindblad RW, et al. Oral Immunotherapy for Treatment of Egg Allergy in Children. *N Engl J Med*. 2012 Jul 19;367(3):233–43.
 19. Gamboa PM, Garcia-Lirio E, Gonzalez C, Gonzalez A, Martinez-Aranguren R M SML. Is the Quantification of Antigen-Specific Basophil Activation a Useful Tool for Monitoring Oral Tolerance Induction in Children With Egg Allergy? *J Investig Allergol Clin Immunol*. 2016 Feb 1;26(1):25–30.
 20. De Vlieger L, Nuyttens L, Matton C, Diels M, Verelst S, Leus J, et al. Guided Gradual Egg-Tolerance Induction in Hen's Egg Allergic Children Tolerating Baked Egg: A Prospective Randomized Trial. *Front Allergy*. 2022;3(May):1–14.
 21. Bartnikas LM, Sheehan WJ, Larabee KS, Petty C, Schneider LC, Phipatanakul W. Ovomucoid Is Not Superior to Egg White Testing in Predicting Tolerance to Baked Egg. *J Allergy Clin Immunol Pract*. 2013 Jul;1(4):354-360.e2.
 22. Ando H, Movérare R, Kondo Y, Tsuge I, Tanaka A, Borres MP, et al. Utility of ovomucoid-specific IgE concentrations in predicting symptomatic egg allergy. *J Allergy Clin Immunol*. 2008;122(3):583–8.
 23. Santos AF, Douiri A, Bécares N, Wu SY, Stephens A, Radulovic S, et al. Basophil activation test discriminates between allergy and tolerance in peanut-sensitized children. *J Allergy Clin Immunol*. 2014 Sep;134(3):645–52.
 24. Lyons JJ, Sun G, Stone KD, Nelson C, Wisch L, O'Brien M, et al. Mendelian inheritance of elevated serum tryptase associated with atopy and connective tissue abnormalities. *J Allergy Clin Immunol*. 2014 May;133(5):1471–4.

25. Claude M, Lupi R, Picariello G, Drouet M, Larré C, Denery-Papini S, et al. Digestion differently affects the ability of native and thermally aggregated ovalbumin to trigger basophil activation. *Food Res Int*. 2019 Apr;118(September 2017):108–14.
26. Benedé S, López-Expósito I, Molina E, López-Fandiño R. Egg proteins as allergens and the effects of the food matrix and processing. *Food Funct*. 2015;6(3):694–713.
27. Nilsson SF, Lilja G, Järnbert-Pettersson H, Alm J. Relevance of low specific IgE levels to egg, milk and peanut in infancy. *Clin Exp Allergy*. 2019 Mar;49(3):308–16.
28. Foong R, Santos AF. Biomarkers of diagnosis and resolution of food allergy. *Pediatr Allergy Immunol*. 2021 Feb;32(2):223–33.
29. Calvani M, Bianchi A, Reginelli C, Peresso M, Testa A. Oral Food Challenge. *Medicina (B Aires)*. 2019 Sep 27;55(10):651.
30. Schoos AM, Bullens D, Chawes BL, Costa J, De Vlieger L, DunnGalvin A, et al. Immunological Outcomes of Allergen-Specific Immunotherapy in Food Allergy. *Front Immunol*. 2020;11(November).
31. De Week AL, Sanz ML, Gamboa PM, Aberer W, Bienvenu J, Blanca M, et al. Diagnostic tests based on human basophils: more potentials and perspectives than pitfalls. II. Technical issues. *J Investig Allergol Clin Immunol*. 2008;18(3):143–55.
32. Chinthrajah RS, Jones SM, Kim EH, Sicherer SH, Shreffler W, Lanser BJ, et al. Updating the CoFAR Grading Scale for Systemic Allergic Reactions in Food Allergy. *J Allergy Clin Immunol*. 2022;1–6.

LEGEND TO FIGURES

Figure 1. 8-month gradual egg-introduction protocol. OFC: oral food challenge, mo: month. Blood samples were drawn prior to and 1 hour after each OFC to evaluate egg sIgE, tryptase level and complement activation. During the home introduction, age-appropriate portions were incorporated 2-3 times/week into the child's diet. Parents monitored the frequency of consumption, allergic reactions, medication or illness by food diary. The child could proceed to the next OFC if no allergic reactions occurred during the introduction of the previous egg preparation at home (*). Adverse allergic events were categorized according to the CoFAR grading scale for allergic reactions (version 3.0) [32]. The amount of egg white protein indicated equals the cumulative dose administered during the OFC (Table S1).




Figure 2. Cubic spline regression analysis of the basophil CD63 dose-response to increasing concentrations of cake (A), hard-boiled egg (B), omelet (C), soft-boiled egg (D), and raw egg (E) in egg allergic (EA, red), complete egg tolerant (ET, green) and non-egg-sensitized non-allergic (NEA, brown) children. The regression curves are shown in bold and represent the mean %CD63⁺ basophils for each group. The shaded area represents the 95% confidence interval. EC50 values were obtained from the non-linear regression analysis for EA and ET children.




Figure 3. Cubic spline regression analysis of basophil CD63 dose-response to increasing concentrations of cake (A), hard-boiled egg (B), omelet (C), soft-boiled egg (D), and raw egg (E) during the first (V1) and last (V5) study visit of the pre-TET1-II study. Data of all participating children were included, excluding non-responders (n=2). The regression curves are shown in bold and represent the mean %CD63⁺ basophils for each group. The shaded area represents the 95% confidence interval. EC50 values were obtained from the non-linear regression analysis for V1 and V5.





Figure 4. Basophil CD63 dose-response to cake (A), hard-boiled egg (B), omelet (C), soft-boiled egg (D), and raw egg (E) during the first and last study visit of the pre-TETI-II study (paired data). Data of all participating children were included, excluding non-responders (n=2). The blue line represents the defined cutoff for each concentration (Table 2).



TABLES

Table 1. Characteristics of the egg allergic and complete egg tolerant children at the time of performing the basophil activation test.

	Age (y)	Gender	Initial egg allergic reaction*	Total IgE (kU/L)	IgE EW (kUA/L)	IgE EY (kUA/L)	IgE OVM (kUA/L)	IgE OVA (kUA/L)
EA 1	4	Girl	Grade III	703	12.1	5.89	7.96	8.96
EA 2	2	Boy	Grade I	864	3.02	0.90	2.04	1.00
EA 3	3	Boy	Grade I	297	4.02	1.19	3.8	1.02
EA 4	4	Boy	Grade I	939	14.9	17.2	4.44	5.31
EA 5 ⁺	<1	Girl	Grade II	256	16.3	4.77	8.42	12.3
EA 6	2	Girl	Grade I	1467	32	5.89	39.4	6.03
EA 7	2	Girl	Grade II	187	13	1.7	20.8	2.31
EA 8	3	Boy	Grade I	280	29	6.86	29.6	12.4
EA 9	1	Girl	Grade I	19	1.15	0.32	1.08	0.62
EA 10	6	Boy	Grade I	2018	5.43	2.16	4.46	3.31
MED (IQR)	2.5 (1.75-4)			500 (238.8-1071)	12.55 (3.77-19.48)	4.77 (1.7-5.89)	6.21 (3.36-23)	5.31 (1.67-10.63)
ET 1	8	Boy	Grade I	278	0.82	0.1	0.64	0.1
ET 2	2	Boy	Grade I	23	0.77	0.24	0.1	1
ET 3 ⁺	5	Girl	Grade I	105	0.23	0.1	0.33	0.1
ET 4	2	Boy	Grade I	8	0.1	0.1	0.1	0.1
ET 5	10	Boy	Grade I	2814	1.2	0.91	0.49	1.36
ET 6	8	Boy	Grade I	2133	0.16	0.12	0.16	0.13
ET 7	8	Boy	Grade I	1232	0.4	0.13	0.52	0.15
ET 8	16	Girl	Grade II	120	0.11	-	0.1	0.12
ET 9	4	Girl	Grade I	51	0.32	0.1	0.25	-
ET 10	5	Boy	Grade I	58	6.13	1.66	0.68	3.62
MED (IQR)	6.5 (3.5-8.5)			112.5 (44-1457)	0.36 (0.15-0.92)	0.10 (0.1-0.91)	0.25 (0.10-0.51)	0.11 (0.10-1.66)

EA: egg allergic, ET: complete egg tolerant, EW: egg white, EY: egg yellow, OVM: ovomucoid, OVA: ovalbumin. ⁺ Non-responder to anti-IgE. * Based on medical record, according to the CoFAR grading scale for allergic reactions (version 3.0) [32].

Table 2. Optimal cutoff values for %CD63⁺ basophils to the five egg extracts with the largest area under the ROC curve.

Extract	Concentration	AUC ROC	Cutoff (%CD63 ⁺ basophils)	Sensitivity (%)	Specificity (%)
Cake	100 µg/mL	1 (1-1)	14.40	100 (84.54-100)	100 (70.09-100)
	<u>10 µg/mL</u>	1 (1-1)	<u>12.14</u>	100 (84.54-100)	100 (70.09-100)
	1 µg/mL	1 (1-1)	2.14	100 (84.54-100)	100 (70.09-100)
	0.1 µg/mL	0.65 (0.41-0.89)	0.52	66.67 (45.37-82.81)	55.56 (26.67-81.12)
Hard-boiled egg	100 µg/mL	0.98 (0.96-1)	7.27	95.24 (77.33-99.76)	100 (70.09-100)
	<u>10 µg/mL</u>	1 (1-1)	<u>10.75</u>	100 (84.54-100)	100 (70.09-100)
	1 µg/mL	1 (1-1)	9.59	100 (84.54-100)	100 (70.09-100)
	0.1 µg/mL	1 (1-1)	4.09	100 (84.54-100)	100 (70.09-100)
Omelet	100 µg/mL	1 (1-1)	13.94	100 (84.54-100)	100 (70.09-100)
	<u>10 µg/mL</u>	1 (1-1)	<u>13.94</u>	100 (84.54-100)	100 (70.09-100)
	1 µg/mL	1 (1-1)	14.26	100 (84.54-100)	100 (70.09-100)
	0.1 µg/mL	1 (1-1)	3.27	100 (84.54-100)	100 (70.09-100)
Soft-boiled egg	100 µg/mL	0.99 (0.96-1)	10.21	95.24 (77.33-99.76)	100 (70.09-100)
	<u>10 µg/mL</u>	1 (1-1)	<u>13.88</u>	100 (84.54-100)	100 (70.09-100)
	1 µg/mL	1 (1-1)	10.27	100 (84.54-100)	100 (70.09-100)
	0.1 µg/mL	0.98 (0.94-1)	2.30	90.48 (71.09-98.31)	100 (70.09-100)
Raw egg	100 µg/mL	1 (1-1)	16.44	100 (84.54-100)	100 (70.09-100)
	<u>10 µg/mL</u>	1 (1-1)	<u>11.68</u>	100 (84.54-100)	100 (70.09-100)
	1 µg/mL	1 (1-1)	11.93	100 (84.54-100)	100 (70.09-100)
	0.1 µg/mL	0.99 (0.98-1)	1.74	95.24 (77.33-99.76)	100 (70.09-100)

AUC ROC: area under the receiver operating characteristic curve. Sensitivity and specificity are expressed as means (95% confidence interval). The underlined concentration and cutoff were applied in the pre-TETI-II study.

Table 3. Baseline demographic and clinical characteristics of patients included in the pre-TETI-II study.

	Age (y)	Gender	Atopy	Initial egg allergic reaction*	Total IgE (kU/L)	IgE EW (kUA/L)	IgE EY (kUA/L)	IgE OVM (kUA/L)	IgE OVA (kUA/L)
L1	7	Boy	AE, AA, AR	Grade I	487	0.48	0.12	0.52	0.12
L2	9	Boy	AE, AA, AR	Grade III	2961	5.39	4.17	0.68	8.06
L3	4	Boy	AE	Grade III	96	0.24	0.1	0.1	0.32
L4	2	Girl	AE	Grade I	207	0.52	0.17	0.1	0.56
L5	2	Girl	AE	Grade III	686	0.24	0.12	0.1	0.26
L6	2	Boy	AE	Grade I	369	0.89	0.56	0.1	1.11
L7	2	Boy	AE	Grade I	64	2	1.18	0.1	1.13
L8	10	Boy	AE, AA	Grade III	576	4.01	1.54	0.8	3.77
L9	2	Girl	AE	Grade III	929	0.24	0.2	0.1	0.22
L10	8	Girl	AE, AA, AR	Grade III	509	1.25	0.53	1.15	0.64
L11	4	Boy	N/A	Grade III	244	1.44	1.3	0.1	2.02
L12	4	Boy	AE, AR	Grade I	1511	2.74	1.45	2.19	1.69
MED (IQR)	4 (2-7.75)				498 (216.3-868.3)	1.070 (0.30-2.56)	0.55 (0.13-1.41)	0.10 (0.10-0.77)	0.88 (0.28-1.94)

AE: atopic eczema, AA: allergic asthma, AR: allergic rhinitis, EW: egg white, EY: egg yellow, OVM: ovomucoid, OVA: ovalbumin. * Based on medical record, according to the CoFAR grading scale for allergic reactions (version 3.0) [32].

Table 4. Basophil response (%CD63⁺ basophils) to 10 µg/mL of the five egg extracts compared with their corresponding oral food challenge during the pre-TETI-II study.

	Cake V1	Hard-boiled egg V2	Omelet V3	Soft-boiled egg V4	Raw egg V5
L1			6.21		3.75
L2		37.96			
L3		1.11		2.57	1.34
L4	0.11	5.64	6.26	3.76	0.09
L5		0	0	0	0.25
L6	7.14	1.15	2.9	6.16	0
L7	12.96	13.96	22.76	14.91	5.78
L8	9.49	0.61			
L9	0.09	0.24	0	0.51	0
L10	0	1.36	0.85	1.92	0
L11	1.33	0	2.45	9.94	5.29
L12		3.21		10.5	10.81
Cutoff	12.14	10.75	13.94	13.88	11.68

Red boxes: children that experienced an allergic reaction during an oral food challenge. Yellow areas: non-responder basophils (L11 became a responder at V3). Green areas: concordant BAT and OFC results. Blue areas: %CD63⁺ basophils above previously defined optimal cutoff. White areas: BAT was not performed due to practical difficulties, or results were not available due to a technical error (L5 cake V1).