

**Supplementary Table I. Cause of anaphylaxis reaction.**

<i>Anaphylaxis (N, %)</i>	<i>110 (20.8)</i>
<b>Food</b>	<b>48 (43.6)</b>
fruits and vegetables	16(14.5)
crustaceans	14 (12.7)
tree nuts	6 (5.5)
anisakis	4 (3.6)
peanuts	4 (3.6)
milk	2 (1.8)
mustard	2 (1.8)
<b>Drugs</b>	<b>46 (41.8)</b>
NSAIDs	32 (29.1)
β-Lactam and	10 (9.09)
quinolones	4 (3.6)
<b>Idiopathic</b>	<b>8 (7.3)</b>
<b>Hymenoptera Sting</b>	<b>2 (1.8)</b>
<b>Latex</b>	<b>2 (1.8)</b>
<b>Exercise induce allergic reaction</b>	<b>2 (1.8)</b>
<b>Mastocytosis</b>	<b>2 (1.8)</b>

**Supplementary Table II. Treatment by primary care physician**

<b>Treatment by primary care physician</b>	<b>(N, %)</b>
<b>In-Patient</b>	<b>110</b>
Epinephrine	<b>20 (18.2)</b>
Antihistamine	<b>82 (74.5)</b>
Systemic corticosteroid	<b>88 (80)</b>
<b>At discharge</b>	
Auto-injectable epinephrine	<b>8 (7.2)</b>
Antihistamine	<b>90 (81.8)</b>
Systemic corticosteroid	<b>48 (43.2)</b>