

## Asthma App Use and Interest Among Patients With Asthma: A Multicenter Study

### SUPPLEMENTARY MATERIAL

Supplementary Table S1. Participants characteristics (n=336).

	Total (n=336)	Adolescents (n=104)	Adults (n=232)
Age y, mean(SD)	34.3 (19.2)	14.7 (1.4)	43.2 (16.8)
Female	207 (62%)	43 (41%)	164 (71%)
Civil Status <sup>a</sup>			
Singled	182 (54%)	104 (100%)	78 (34%)
Married/ Civil union	124 (37%)	0	124 (53%)
Divorced	20 (6%)	0	20 (9%)
Widowed	9 (3%)	0	9 (4%)
Education <sup>b</sup>			
Primary school or below	40 (12%)	3 (3%)	37 (16%)
Elementary school	125 (37%)	76 (73%)	49 (21%)
High school	95 (28%)	23 (22%)	72 (31%)
University	70 (21%)	0	70 (30%)
Occupation <sup>c</sup>			
Employed	138 (41%)	0	138 (60%)
Student	133 (40%)	104 (100%)	30 (13%)
Retired	34 (10%)	0	34 (15%)
Unemployed / Not working due to poor health	18 (5%)	0	18 (8%)
Other	9 (3%)	0	9 (4%)
BMI Kg/m <sup>2</sup> , mean (SD)	24.8 (5.6)	21.4 (4.1)	26.4 (5.5)
Smoking status <sup>d</sup>			
Never smoker	256 (76%)	96 (92%)	160 (69%)
Ex-smoker	55 (16%)	6 (6%)	49 (21%)
Current smoker	23 (7%)	2 (2%)	21 (9%)
GINA assessment symptom control <sup>e</sup>			
Well controlled	175 (52%)	48 (46%)	127 (55%)
Partly controlled	97 (29%)	40 (39%)	57 (25%)
Uncontrolled	61 (18%)	16 (15%)	45 (20%)

Values are shown as n(%) unless otherwise indicated. Abbreviations: BMI, body mass index; GINA, Global Initiative for Asthma. a – 1 missing value; b – 6 missing values; c – 3 missing values; d – 2 missing values; e – 3 missing values.