

SUPPLEMENTARY MATERIAL

Table. Epidemiological and clinical characteristics, skin prick test results, and specific IgE (Siemens Immulite 2000/Xpi®) to whiff and other fish species:

Patient	Gender	Age	Symptoms	Skin prick test*/ sIgE (kU _A /L)**					
				Cod	Tuna	Salmon	Hake	Whiff	Sole
1	F	2	U	-/-	-/-	-/-	-/-	+/1.4	-/-
2	F	7	U	-/-	-/-	-/-	-/-	+/ 0.4	-/-
3	M	1	A	-/-	-/-	-/-	/-	+/1.4	-/-
4	F	21	U/An	-/-	-/-	-/-	-/-	+/0.5	-/-
5	M	4	SAO	-/-	-/-	-/-	-/ 0.7	+/2.1	-/-
6	M	12	U/An	-/1.4	-/-	-/-	-/ -	+/0.4	-/0.9
7	M	52	A	-/0.7	-/-	-/-	-/ 1.6	+/10.5	-/1
8	M	2	A	-/1.5	-/-	-/2.2	-/ 0.9	+/10.3	-/2.5
9	F	1	U	ND/1.7	-/0.5	-/0.9	ND/ 1.7	+/2.6	ND/1.1
10	F	25	U/An	-/-	-/-	-/-	-/ -	+/0.9	-/-

P: Patient; Gender: M (male), F (female); Age (years): allergic reaction age of onset; Symptoms: A (anaphylaxis), U (urticaria), An (angioedema), OAS (oral allergy syndrome).

*+: Positive. Wheal diameters larger than 3 mm were considered to indicate a positive result; -: Negative; ND: not done. **Positive values were those ≥ 0.35 kUA/L