

## SUPPLEMENTARY MATERIAL

**Table S1.** Characteristics of the study patients after matching by age and gender

	Total (n = 30)	Patients with SLH (n = 15)	Patients without SLH (n = 15)	p value
<b>Female, N (%)</b>	12(40%)	6(40%)	6(40%)	1.000
<b>Age (year)</b>	57.5(50.8-68.0)	57.0(52.0-68.0)	58.0(50.0-69.0)	0.967
<b>BMI (kg/m<sup>2</sup>)</b>	25.3(23.4-27.7)	25.2(23.1-27.9)	25.6(24.1-27.7)	0.999
<b>Duration of asthma (years)</b>	17.0(6.5-27.8)	12.0(5.0-26.0)	20.0(9.0-32.0)	0.361
<b>Late-onset asthma, N (%)</b>	12(40%)	5(33.3%)	7(46.7%)	0.456
<b>ACT score</b>	21.0(17.8-24.0)	22.0(19.0-24.0)	20.0(17.0-23.0)	0.142
<b>Medication</b>				
Medium or high dose				
ICS/LABA/LAMA, N (%)	23(76.7%)	12(80.0%)	11(73.3%)	0.666
OCS, N (%)	13(43.3%)	5(33.3%)	8(53.3%)	0.268
Biologics, N (%)	14(46.7%)	7(46.7%)	7(46.7%)	1.000
<b>FeNO (ppb)</b>	34.0(19.3-80.5)	54.0(17.0-101.0)	20.0(29.0-42.0)	0.329
<b>Blood eosinophil (/µL)</b>	241.5(141.8-317.3)	246.0(138.0-359.0)	237.0(143.0-304.0)	0.917
<b>PEFR (L/min)</b>	300(255.0-410.0)	300.0(240.0-380.0)	310.0(260.0-440.0)	0.486
<b>Spirometry</b>				
FEV <sub>1</sub> /FVC (%)	65.0(53.8-73.0)	65.7(52.3-75.2)	64.2(54.3-71.6)	0.885
FEV <sub>1</sub> (% of predicted)	70.6(60.5-88.4)	63.5(60.2-87.8)	84.0(60.6-100.6)	0.254
FVC (% of predicted)	87.4(73.9-95.1)	82.2(66.1-88.7)	91.4(85.6-107.8)	<b>0.044</b>
FEF <sub>25-75</sub> (% of predicted)	39.5(21.8-54.2)	37.0(20.0-53.3)	44.0(22.4-68.0)	0.604
BDR (+), N (%)	9(30.0%)	4(13.3%)	5(16.7%)	0.690
TLC (% of predicted)	103.0(94.8-116.3)	105.0(98.0-118.0)	98.0(94.0-111.0)	0.309
RV/TLC	40.5(38.0-50.0)	50.0(45.0-53.0)	38.0(36.0-39.0)	<b>0.000</b>

---

**Impulse oscillometry**

R <sub>5</sub> [kPa/(L/s)]	0.44(0.39-0.54)	0.48(0.39-0.55)	0.43(0.31-0.49)	0.191
R <sub>5</sub> (% of predicted)	160.8(131.2-219.9)	178.5(138.7-235.1)	142.8(129.2-198.8)	0.221
R <sub>5</sub> -R <sub>20</sub> [kPa/(L/s)]	0.13(0.10-0.18)	0.14(0.10-0.19)	0.12(0.08-0.16)	0.197
R <sub>5</sub> -R <sub>20</sub> (% of predicted)	224.7(161.6-428.8)	277.5(184.3-481.7)	208.3(149.2-360.4)	0.330
F <sub>res</sub> (Hz)	18.5(15.7-22.2)	18.9(16.9-24.7)	16.9(14.4-21.6)	0.158
F <sub>res</sub> (% of predicted)	146.6(128.4-185.8)	151.2(132.3-192.8)	142.9(122.5-183.5)	0.290
X <sub>5</sub> [kPa/(L/s)]	-0.2(-0.32 - -0.11)	-0.27(-0.36- -0.20)	-0.14(-0.20 -0.09)	<b>0.007</b>
X <sub>5</sub> (% of predicted)	187.0(112.9-304.5)	291.0(182.6-348.8)	128.9(94.6-225.9)	<b>0.010</b>
AX (kPa/L)	1.42(0.70-2.03)	1.64(1.05-2.04)	0.96(0.38-1.53)	<b>0.049</b>
AX (% of predicted)	445.8(217.1-877.1)	518.4(266.4-900.4)	292.0(152.4-774.6)	0.101

---

SLH, static lung hyperinflation, defined as the ratio of residual volume (RV) to total lung capacity (TLC) ratio more than 0.4; BMI, body mass index; Late-onset asthma, defined as age of asthma onset  $\geq$  40 years old; ACT, asthma control test; ICS/LABA/LAMA, inhaled corticosteroid, long-acting beta-2 agonist and long-acting muscarinic antagonist combination therapy; OCS, oral corticosteroid; FeNO, fractional exhaled nitric oxide; PEFR: peak expiratory flow rate; FEV<sub>1</sub>: forced expiratory volume in the first second; FVC: forced vital capacity; FEF<sub>25-75%</sub> : forced expiratory flow between 25 and 75% of forced vital capacity; BDR (+), a positive bronchodilator response, defined as an increase in either FEV<sub>1</sub> or FVC for more than 12% and 200 mL from the baseline in response to a short-acting beta-2 agonist; R<sub>5</sub>, resistance in 5 Hz; R<sub>20</sub>, resistance in 20 Hz; F<sub>res</sub>, resonant frequency; X<sub>5</sub>, reactance in 5 Hz; AX, area under reactance curve between 5 Hz and resonant frequency; data are shown as number (%) for categorical variables and median (interquartile range, IQR) for non-normally distributed variables. *p* values were calculated by the Mann-Whitney *U* test and the values <0.05 were considered statistically significant.

**Table S2.** Performance of different IOS parameters to detect SLH in patients with severe asthma

Variables	Cutoff	Sensitivity	Specificity	LR (+)	LR (-)	AUC	Youden	p value
	value	(%)	(%)			(95%CI)	Index	
R <sub>5</sub> [kPa/(L/s)]	>0.49	67.1	87.5	5.37	0.38	0.76 (0.67-0.84)	0.55	<0.0001
R <sub>5</sub> (% of predicted)	≤267.6	96.4	20.8	1.22	0.17	0.53 (0.43-0.62)	0.17	0.7313
R <sub>20</sub> [kPa/(L/s)]	>0.34	51.2	83.3	3.07	0.59	0.70 (0.60-0.78)	0.35	0.0009
R <sub>20</sub> (% of predicted)	>141.3	57.3	66.7	1.72	0.64	0.63 (0.53-0.72)	0.24	0.0359

IOS, impulse oscillometry; SLH, static lung hyperinflation, defined as the ratio of residual volume (RV) to total lung capacity (TLC) ratio more than 0.4; R<sub>5</sub>, resistance in 5 Hz; R<sub>20</sub>, resistance in 20 Hz; LR (+), positive likelihood ratio; LR (-), negative likelihood ratio; AUC: area under the receiver operating characteristic curve. 95% CI: 95% of confidence interval.